

Citation for published version:

Rousis, NI, Gracia-Lor, E, Zuccato, E, Bade, R, Baz Lomba, JA, Castrignanò, E, Causanilles, A, Covaci, A, de Voogt, P, Hernández, F, Kasprzyk-Hordern, B, Kinyua, J, McCall, AK, Plósz, BG, Ramin, P, Ryu, Y, Thomas, KV, van Nuijs, A, Yang, Z & Castiglioni, S 2017, 'Wastewater-based epidemiology to assess pan-European pesticide exposure', *Water Research*, vol. 121, pp. 270-279. <https://doi.org/10.1016/j.watres.2017.05.044>

DOI:

[10.1016/j.watres.2017.05.044](https://doi.org/10.1016/j.watres.2017.05.044)

Publication date:

2017

Document Version

Peer reviewed version

[Link to publication](#)

Publisher Rights

CC BY-NC-ND

University of Bath

Alternative formats


If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



Wastewater-based epidemiology to assess pan-European pesticide exposure

Nikolaos I. Rousis^{a*}, Emma Gracia-Lor^{a,b}, Ettore Zuccato^a, Richard Bade^{b,m}, Jose Antonio Baz-Lomba^c, Erika Castrignanò^d, Ana Causanilles^e, Adrian Covaci^f, Pim de Voogt^e, Félix Hernández^b, Barbara Kasprzyk-Hordern^d, Juliet Kinyua^f, Ann-Kathrin McCall^g, Benedek Gy. Plósz^{h,i}, Pedram Ramin^{h,j}, Yeonsuk Ryu^c, Kevin V. Thomas^{c,k}, Alexander van Nuijs^f, Zhugen Yang^{d,l}, Sara Castiglioni^{a*}

^a IRCCS — Istituto di Ricerche Farmacologiche “Mario Negri”, Department of Environmental Health Sciences, Via La Masa 19, 20156 Milan, Italy

^b Research Institute for Pesticides and Water, University Jaume I, Avda. Sos Baynat s/n, E-12071 Castellón, Spain

^c Norwegian Institute for Water Research (NIVA), Gaustadalléen 21, 0349 Oslo, Norway

^d University of Bath, Department of Chemistry, Faculty of Science, Bath BA2 7AY, United Kingdom

^e KWR Watercycle Research Institute, Chemical Water Quality and Health, P.O. Box 1072, 3430 BB, Nieuwegein, The Netherlands and IBED-University of Amsterdam, The Netherlands.

^f Toxicological Center, Department of Pharmaceutical Sciences, Campus Drie Eiken, University of Antwerp, Universiteitsplein 1, 2610 Wilrijk-Antwerp, Belgium

^g Eawag, Swiss Federal Institute of Aquatic Science and Technology, CH-8600, Dübendorf, Switzerland

^h Department of Environmental Engineering, Technical University of Denmark, Bygningstorvet,
Building 115, DK-2800 Kgs. Lyngby, Denmark

ⁱ Department of Chemical Engineering, University of Bath, Claverton Down, Bath BA2 7AY, UK

^j Department of Chemical and Biochemical Engineering, Technical University of Denmark, Søltøfts
Plads, Building 229, DK-2800 Kgs. Lyngby, Denmark

^k Queensland Alliance for Environmental Health Science (QAEHS), University of Queensland, 39
Kessels Road, Coopers Plains QLD 4108, Australia

^l Division of Biomedical Engineering, School of Engineering, University of Glasgow, Glasgow G12
8LT United Kingdom

^m School of Pharmacy and Medical Sciences, University of South Australia, Adelaide, South Australia 5000,
Australia

*Corresponding authors: Sara Castiglioni (sara.castiglioni@marionegri.it); Nikolaos I. Rousis
(nrousis@chem.uoa.gr)

Abstract

Human biomonitoring, i.e. the determination of chemicals and/or their metabolites in human specimens, is the most common and potent tool for assessing human exposure to pesticides, but it suffers from limitations such as high costs and biases in sampling. Wastewater-based epidemiology (WBE) is an innovative approach based on the chemical analysis of specific human metabolic excretion products (biomarkers) in wastewater, and provides objective and real-time information on xenobiotics directly or indirectly ingested by a population. This study applied the WBE approach for the first time to evaluate human exposure to pesticides in eight cities across Europe. 24h-composite wastewater samples were collected from the main wastewater treatment plants and analyzed for urinary metabolites of three classes of pesticides, namely triazines, organophosphates and pyrethroids, by liquid chromatography-tandem mass spectrometry. The mass loads (mg/day/1000 inhabitants) were highest for organophosphates and lowest for triazines. Different patterns were observed among the cities and for the various classes of pesticides. Population weighted loads of specific biomarkers indicated higher exposure in Castellon, Milan, Copenhagen and Bristol for pyrethroids, and in Castellon, Bristol and Zurich for organophosphates. The lowest mass loads (mg/day/1000 inhabitants) were found in Utrecht and Oslo. These results were in agreement with several national statistics related to pesticides exposure such as pesticides sales. The daily intake of pyrethroids was estimated in each city and it was found to exceed the acceptable daily intake (ADI) only in one city (Castellon, Spain). This was the first large-scale application of WBE to monitor population exposure to pesticides. The results indicated that WBE can give new information about the “average exposure” of the population to pesticides, and is a useful complementary biomonitoring tool to study population-wide exposure to pesticides.

Keywords: Urban wastewater; Mass spectrometry; Pesticides; Human urinary metabolites; Biomonitoring; Human intake

62

63 **1 Introduction**

64 Pesticides play an important role in agriculture by protecting plants and plant products
65 against harmful organisms and their action, and helping boost the growth of crops. Meeting the
66 demand in food supply will be one of the great challenges in the near future, since the global
67 population is expected to grow to nine billion by the middle of the century (Godfray et al., 2010). In
68 order to raise food production, an increased pesticides use is expected. Taking into account that
69 thousands of tons of pesticides are yearly applied in agriculture, homes, gardens, sports fields, and
70 public areas (Grube et al., 2011), contamination of the environment most likely will further increase
71 and human exposure to pesticides will continue being a matter of substantial concern in the near
72 future.

73 Many “old and harmful” pesticides, such as p,p-dichlorodiphenyl-trichloroethane (p,p’-
74 DDT), have been banned because of their toxicity and they were replaced by less-persistent
75 pesticides, such as organophosphates and pyrethroids (Barr, 2008; López et al., 2005). Pesticides
76 provide mankind with many benefits, but at the same time have the potential to pose risks for
77 human health due to widespread use and high biological activity (Cooper and Dobson, 2007). For
78 instance, pesticides exposure has positive association with the development of idiopathic
79 Parkinson’s disease, neurobehavioral and neuropsychological disorders, respiratory symptoms or
80 diseases, and sperm DNA damage (Allen and Levy, 2013; Mamane et al., 2015; Saillenfait et al.,
81 2015; Stallones and Beseler, 2016). However, in the last two decades, the concept of “green
82 chemistry” has been promoted and the agrochemical industry has focused on less toxic substances
83 (Garrison, 2004).

84 The general population is exposed to pesticides mainly through diet and household use
85 (Aprea, 2012). Human biomonitoring (HBM) is the main tool for assessing exposure and consists in

the measurement of chemicals and/or their metabolites in body fluids or tissues (Barr, 2008; Yusa et al., 2015). The reliability of HBM depends on the selection of a proper biomarker that reflects the exposure to the parent compound, and is specific and detectable in the investigated matrices. Urine is the preferred human biological matrix, since it is easy to collect and non-invasive and it is also accessible in large volumes allowing the determination of very low concentrations of chemicals compared to other fluids (Wessels et al., 2003). Extensive HBM studies have analyzed the urine of thousands of individuals to investigate pesticide exposure in the general population (Barr et al., 2010, 2004; Heudorf and Angerer, 2001; McKelvey et al., 2013; Ye et al., 2015). Despite their power to evaluate exposure to chemicals, HBM studies suffer by limitations such as high costs for sample collection and analysis, ethical issues and data analysis to extrapolate individual results to the whole population. Moreover, urine sampling can reflect only a momentary snapshot of exposure due to sampling procedures (i.e. morning urine collection), and excretion profiles may vary throughout the day/days because of the short half-lives in the human body of most of pesticides.

Wastewater-based epidemiology (WBE) is a recent approach for the retrieval of epidemiological information from wastewater through the analysis of specific human metabolic excretion products (biomarkers) (Castiglioni et al., 2014). It can be described as a collective urine test, as the wastewater from a city pools the anonymous urine samples of thousands of individuals. WBE was originally developed in Italy to estimate illicit drug consumption in a population (Zuccato et al., 2008) and has later been applied worldwide with promising results (Banta-Green et al., 2009; Ort et al., 2014). New possibilities permit information on public health and lifestyles (Thomas and Reid, 2011; Venkatesan and Halden 2014). The main advantage of WBE is to provide objective, real-time information on substances directly or indirectly ingested daily by a population, with a clear potential to provide complementary data for epidemiological studies and to overcome some of the HBM limitations.

The first exploratory study proposing WBE as a novel biomonitoring tool to evaluate the exposure of the general population to pesticides was recently performed (Rousis et al., 2016). Several metabolites of organophosphates, triazines and pyrethroids were detected in raw wastewater and their frequency of detection and abundance were in agreement with the profiles reported in urine of HBM studies (Rousis et al., 2016). Later three human urinary metabolites of pyrethroids were selected and used to back-calculate the population intake of pyrethroids in Italy (Rousis et al., 2017). This study indicated for the first time that WBE can be employed as a complementary biomonitoring tool to the HBM studies, but more data and a wider scale of investigation were necessary in order to confirm these preliminary results.

The aim of the present study was to apply for the first time this new WBE approach in eight countries across Europe and to evaluate the pan-European human exposure to pesticides in order to validate the method by comparing results with international statistics. 24-h composite raw wastewater samples were collected and analyzed for organophosphate, triazine and pyrethroid metabolites. The results for the cities were compared and population-wide pyrethroid intake was estimated. To the best of our knowledge, this is the first WBE study designed to assess human exposure to pesticides at a European scale.

2 Materials and methods

2.1 Chemicals and reagents

Hydrochloric acid (HCl, 37%) and acetonitrile for liquid chromatography-mass spectrometry (LC-MS) were purchased from Riedel de Haen (Seelze, Germany); methanol (MeOH) for pesticide analysis from Carlo Erba Reagents (Italy); triethylamine and acetic acid from Fluka (Buchs, Switzerland). HPLC grade Milli-Q water was obtained with a Milli-RO Plus 90 apparatus (Millipore, Molsheim, France). Analytical standards for diethyl phosphate (DEP, purity 97.6%),

chlorpyrifos (CPF, purity 99.9%), chlorpyrifos methyl (CPF-MET, purity 99.5%) and 3,5,6-trichloro-2-pyridinol (TCPY, purity 99.5%) were purchased from Chemical Research 2000 (Rome, Italy). Atrazine (ATZ, purity 97.5%), atrazine desethyl (DEA, purity 99.9%), terbutylazine desethyl (DES, purity 97.4%), atrazine desisopropyl (DIA, purity 95.4%), dimethyl chlorophosphate (DMCIP, purity 96%), dimethyl chlorothiophosphate (DMCITP, purity 97%), and O,O-diethyl thiophosphate (DETP, purity 98%) potassium salt were supplied by Sigma-Aldrich (Schnelldorf, Germany). Atrazine mercapturate (AM, purity 95.0%), 3-(2,2-dichlorovinyl)-2,2-dimethyl-(1-cyclopropane)carboxylic acid (DCCA, purity 99.0%), 3-phenoxybenzoic acid (3-PBA, purity 99.0%), 2-isopropyl-6-methyl-4-pyrimidinol (IMPY, purity 99.5%), cis-3-(2,2-dichlorovinyl)-2,2-dimethyl-(1-cyclopropane) carboxylic acid (cis-DCCA, purity 98%) and malathion monocarboxylic acid (MMA, purity 97.0%) were purchased from Lab Service Analytica (Bologna, Italy). Isotopically labeled compounds (deuterated or ^{13}C -enriched) were used as internal standards (IS). 3-Phenoxybenzoic acid- C_6 (3-PBA- $^{13}\text{C}_6$, phenoxy- $^{13}\text{C}_6$, 99%; purity 98%) and 3,5,6-trichloro-2-pyridinol- C_3 (TCPY- C_3 , 4,5,6- $^{13}\text{C}_3$, 99%; purity 97%) were obtained from Cambridge Isotope Laboratories, Inc. (Massachusetts, USA); atrazine- D_5 (ATZ- D_5 , 99.5%) from Sigma-Aldrich (Schnelldorf, Germany); and chlorpyrifos D_{10} (CPF- D_{10} , 97.0%) from Lab Service Analytica (Bologna, Italy). Dimethyl phosphate (DMP) and dimethyl thiophosphate (DMTP) were synthesized by simple hydrolysis of DMCIP and DMCITP (Hernández et al., 2002; Rousis et al., 2016a).

2.2 Selection of exposure biomarkers

Specific urinary metabolites of pesticides were selected as biomarkers from HBM studies available in literature and official reports of the United States Environmental Protection Agency and the Centers for Disease Control and Prevention, as described elsewhere (Rousis et al., 2016). The biomarkers were chosen according to specific criteria: a) levels in urine; b) frequency of detection;

c) frequency of use of the respective classes of pesticides; d) risks for human health; e) specificity of the metabolites (human excretion *versus* environmental formation).

The selected biomarkers were three parent substances and 15 urinary metabolic products belonging to different pesticide classes. Among triazines, the parent atrazine and the metabolites DES, DIA, DEA and AM were selected. Among pyrethroids 3-PBA, the common metabolite of about 20 synthetic pyrethroids, and *cis*- and *trans*-DCCA, which are the specific metabolites of permethrin, cypermethrin and cyfluthrin were chosen. Among organophosphates, the four alkyl phosphates DEP, DETP, DMP and DMTP, which are common metabolites of a large group of organophosphates, chlorpyrifos, chlorpyrifos methyl and their specific metabolite TCPY, the metabolites of malathion (the α and β isomers of MMA) and the metabolite of diazinon (IMPY) were selected.

The reliability of back-calculation of the exposure to parent chemicals (pesticides) depends strictly on the selection of an appropriate WBE biomarker, which can be either the compound itself or one of its metabolites. Therefore, the selected metabolites were checked to fulfill the requirements of a WBE biomarker, which are: a) measurable in raw wastewater; b) released into sewers only as a result of human excretion; c) a well-defined excretion profile to avoid interference from other exogenous or endogenous sources; d) limited adsorption to suspended matter; e) stable in wastewater during in-sewer transit, sampling and storage (Gracia-Lor et al., 2016). The stability of each compound in wastewater was evaluated through specific laboratory tests (Rousis et al., 2016), and the specificity of each metabolite was assessed by checking the presence of sources other than human metabolism (i.e. any potential environmental transformation) (Rousis et al., 2017 and this study). The results for the selected substances are summarized in Table 1.

2.3 Samples and sampling method

Raw wastewater samples were taken from the inlet of the wastewater treatment plants (WWTPs) of eight European cities: Bristol, the United Kingdom; Brussels, Belgium; Castellon, Spain; Copenhagen, Denmark; Milan, Italy; Oslo, Norway; Utrecht, The Netherlands and Zurich, Switzerland (Figure 1).

Composite 24-h samples of untreated wastewater were collected by automatic sampling devices (Table S1). Sampling was carried out over one week in March 2015. For each WWTP, seven consecutive 24-h samples were collected in high-density polyethylene bottles, transferred to Milan and stored at -20°C until sample treatment.

2.4 Sample pretreatment

The method for sample preparation was published in detail elsewhere (Rousis et al., 2016). Briefly, samples were filtered on a glass microfiber filter GF/A 1.6 μm (Whatman, Kent, U.K.) and on a mixed cellulose membrane filter 0.45 μm (Whatman, Kent, U.K.) before extraction. Solid phase extraction (SPE) was used to extract the target analytes using OASIS[®] HLB 3 cc/60 mg cartridges (Waters Corp., Milford, MA, USA) and an automatic GX-274 ASPEC (Gilson, Middleton, WI, USA) extractor. Samples (50 mL of untreated wastewater) were spiked with 2 ng of a mixture of internal standards and the pH was adjusted to 7.0-7.5, using diluted HCl (12%). Cartridges were conditioned with MeOH (5 mL) and Milli-Q water (3 mL) and samples were passed at a flow rate of 5 mL/min. The cartridges were dried under a nitrogen stream at a flow rate of 10 mL/min for 10 min and eluted with 3 mL of MeOH. Eluates were evaporated under a gentle nitrogen stream at room temperature and dried samples were reconstituted in 100 μL of Milli-Q water and transferred into glass vials for LC-MS/MS analysis.

The alkyl phosphate analytes DEP, DETP, DMP and DMTP were directly injected into the LC-MS/MS system; 500 μL of filtered samples were centrifuged at 2500 rpm for 2 min and 180 μL

of supernatant were collected, spiked with 2 ng of a mixture of internal standards and transferred into glass vials for LC-MS/MS analysis.

2.5 Instrumentation and analytical method

Chromatographic separation was done with an Agilent 1200 Series system (Agilent Technologies, Santa Clara, CA, USA) using an XSELECTTM CSHTM C18 (2.1 × 100 mm, 2.5 μm) column (Waters Corp., Milford, MA, USA). Mass spectrometric analysis done using an AB SCIEX Triple QuadTM 5500 LC-MS/MS System (AB-Sciex, Thornhill, Ontario, Canada). Two or three most abundant product ions of the protonated pseudo-molecular ion of each substance were chosen for analysis which was done both in positive and negative ionization modes using the selected reaction monitoring mode (SRM). Quantification was performed by isotopic dilution. Method limits of detection and quantification are reported in Table S2. The method was fully validated in raw wastewater, as described elsewhere (Rousis et al., 2016).

2.6 Stability of biomarkers and parent pesticides in wastewater

Stability experiments aim to ensure that no degradation of the targeted compounds occurs in the sewage system and during sampling and storage, so no pre-analytical losses occur (McCall et al., 2016). The stability of parent pesticides is crucial, since degradation of these compounds could lead to formation of the targeted biomarker in wastewater, hence to overestimation of human exposure. The stability of metabolites in raw wastewater and the formation of pyrethroid metabolites from the degradation of parent pyrethroids were evaluated in previous studies (Rousis et al., 2016, 2017). The present study investigated the formation of triazine and some organophosphate metabolites after addition of the corresponding parent pesticides in raw wastewater, under different conditions. Parent triazine (atrazine, simazine, propazine, terbutylazine)

and organophosphate pesticides (chlorpyrifos, chlorpyrifos-methyl, malathion, diazinon) were spiked in wastewater to the maximum acceptable concentration ($0.1 \mu\text{g/L}$) for a single pesticide in groundwater, surface water and water intended for human consumption according to EU directives (Commission, 2008, 2006, 1998) to test their stability under controlled conditions (room temperature and 4°C). These temperatures were chosen in order to mimic conditions in the sewer system (room temperature, $\sim 23^\circ\text{C}$; worst case scenario) and during the collection of the composite 24-h samples (occurring at 4°C). Each experiment was run in triplicate and samples were analyzed immediately after spiking (t_0), and after 6 (t_6) and 24 h (t_{24}). Unspiked samples were used as matrix blanks. Analysis of formed DEP, DETP, DMP and DMTP compounds following addition of parent pesticides in wastewater was not performed, since these metabolites are excretion or transformation products of a wide number of pesticides and other substances including flame retardants, plasticizers and industrial chemicals (Rousis et al., 2016).

2.7 Daily mass loads

Daily mass loads of biomarkers were calculated by multiplying the concentrations (ng/L) found in a 24h composite sample of raw wastewater by the daily wastewater flow rate (m^3/day) at the WWTPs (Table S1). Biomarker mass loads (mg/day) were then normalized to the number of people served by each WWTP ($\text{mg/day}/1000$ inhabitants), in order to compare results between different cities.

2.8 Pyrethroid intake and uncertainty evaluation

At present, pyrethroid metabolites (3-PBA and DCCA) were found to be the most suitable biomarkers of exposure according to the specific requirements of WBE (Table 1), so they were used to back-calculate population-wide intake of pyrethroids. Specific correction factors (CFs) were

developed by Rousis et al. (2017) and the following equation was used to estimate pyrethroids intake:

$$P_{YR_{intake}} = \frac{(Conc. \times F) \times CF}{P}$$

where: Conc. is the concentration of each target analyte (ng/L) in wastewater, F is the corresponding flow rate of wastewater in WWTP (m³/day), CF is the specific correction factor for each analyte and P is the population served by each WWTP.

CFs were calculated taking into account the molar mass ratio between parent pesticide and target metabolite and the percentage of excretion of the target metabolite in human urine. Since each metabolite is common to more than one parent substance, the molar mass ratios were calculated using the arithmetic mean of the molecular weights of all parent substances divided by the molecular weight of each metabolite. All human urinary pharmacokinetic studies reporting the excretion rate of metabolites after a dose of the parent substances were considered. The weighted mean (WM) excreted fraction was calculated as the mean percentage of excretion weighted by the number of subjects in each study (Rousis et al., 2017). The following equation was used to calculate CFs:

$$CF = \frac{\frac{Mw \text{ (Parent pesticide)}}{Mw \text{ (metabolite)}}}{WM \text{ excreted fraction (metabolite)}}$$

where: Mw is the molecular weight and WM is the weighted mean of the percentage of excretion of the targeted metabolites.

The procedure used to develop CFs has been described in detail elsewhere (Rousis et al., 2017). CFs were 6.95 for 3-PBA (used to estimate the intake of 20 pyrethroids) and respectively 3.67 and 5.45 for *trans*- and *cis*-DCCA (used to estimate the intake of permethrin, cypermethrin, and cyfluthrin) (Rousis et al., 2017). The intake levels of permethrin, cypermethrin and cyfluthrin (sum of *cis*- and *trans*- levels) estimated by WBE were compared with a toxicological indicator, the

acceptable daily intake (ADI), so as to evaluate the measured levels of exposure in relation to their potential effects on human health.

Uncertainty was evaluated following the available best practice protocols for WBE (Castiglioni et al., 2014, 2013). Sampling procedures were selected to keep uncertainty below 10%, while the analytical procedure was optimized to have an analytical variability lower than 15% (Rousis et al., 2016). The variability of excretion profiles of pyrethroids metabolites was carefully evaluated to assess the uncertainty related to CFs and consequently to the back-calculation. It was calculated as the standard deviation of the percentages of excretion collected from the literature as shown previously (Rousis et al., 2017) and it was lower than 16%. Finally, data normalization to the population served by each WWTP was done considering the most reliable population estimation to keep uncertainty as lower as possible. Nevertheless, as described elsewhere, this is probably the most critical aspect of determining the variability (Castiglioni et al., 2013).

2.9 Data elaboration

Data were analysed using a MultiQuantTM 2.1 software package of Analyst[®] (AB-Sciex, Thornhill, Ontario, Canada). GraphPad Prism (Version 6.0) was used for figures elaboration and statistical analyses which was performed by using an unpaired t-test or a Mann-Whitney test according to the normality of data. All tests were done considering a statistical significance level of $p < 0.05$. Concentrations below the Limit of Quantification (LOQ) were replaced with a value equal to half the LOQ.

3 Results and Discussion

3.1 Stability of metabolites and parent pesticides

The stability experiments showed no formation of triazine and organophosphate metabolites in any of the tested conditions (Table S3). Thus, the percentage variation of the concentration for each metabolite at t_6 and t_{24} respect to t_0 indicated that very small variations occurred for all metabolites. Even though these laboratory experiments were conducted under controlled conditions (pH = 7.0-7.5; room temperature and 4 °C) that are not reproducing the spatial and temporal variability in a sewer system, they can provide indicative information regarding the stability of a compound in wastewater.

3.2 Occurrence of biomarkers in raw wastewater

Concentrations of the biomarkers measured in wastewater are shown in Table 2 with their frequencies of detection. The substances most frequently observed were ATZ and DEA (detection rates 98.2% and 62.5%) among triazines; 3-PBA and *trans*-DCCA (detection rates 98.2% and 96.4%) among pyrethroids; TCPY (detection rate 100%), IMPY (detection rate 87.5%), and DMP and DEP (detection rates 100% and 94.6%) among organophosphates. The other biomarkers had lower frequencies of detection (<40%), and chlorpyrifos, chlorpyrifos-methyl and DMTP were not detected. Mean concentrations ranged from a few ng/L (triazines) to 2.3 µg/L (DMP).

The results were comparable with those of a previous study in seven Italian cities (Rousis et al., 2016). The profiles of the compounds most frequently detected were similar, besides a few exceptions; i.e. the frequency of detection of DES and *cis*-DCCA was higher in Italy (100% and 73%) than in the other European cities (38% and 36%), and CPF was detected in one city in Italy (Rousis et al., 2016), but not in the EU cities (Table 2). The results for the other compounds were quite similar in both studies: AM, CPF-MET and DMTP were not detected; malathion and triazine metabolites were detected sporadically (frequency of detection <40%); and TCPY and DMP were detected in all samples. The highest concentrations in both studies were measured for the alkyl

phosphate metabolites, DEP and DMP, which are metabolic products of most organophosphates, while the triazines group was found at the lowest concentrations (Rousis et al., 2016). The concentrations of *trans*-DCCA were always higher than those of *cis*-DCCA, in accordance with HBM studies, where the *trans*-isomer predominated (Rousis et al., 2017). The *trans*- to *cis*- DCCA ratio is used as an indicator of the route of human exposure and a ratio of 2:1 or higher indicates oral uptake and/or inhalation. This suggests that these metabolites in wastewater resulted mainly from human metabolism, since the ratio was higher than 2:1, as reported previously (Rousis et al., 2017).

3.3 Mass loads of biomarkers in the different cities

The mean mass loads of organophosphates, triazines and pyrethroids (parent and metabolites) expressed as mg/day/1000 inhabitants, are reported in Table S4.

The alkyl phosphates DMP and DEP gave the highest loads (up to 975 mg/day/1000 inh for DMP and 244 mg/day/1000 inh for DEP). These high mass loads were expected, since these substances are metabolic products of most of the organophosphate insecticides used in Europe. These substances also might originate from plasticizers or flame retardants following hydrolysis or from other industrial chemicals (Reemtsma et al., 2011) and are therefore not specific for human exposure. Among the other specific metabolites investigated, the loads of the diazinon metabolite IMPY ranged from 1.3 to 16 mg/day/1000 inh. and the metabolite of chlorpyrifos and chlorpyrifos-methyl, TCPY, ranged from 3.9 to 22 mg/day/1000 inh., suggesting different exposure to these organophosphates in the various countries.

Triazines had the lowest loads, ranging from 0.33 to 5.0 mg/day/1000 inh. Generally, the metabolite mass loads were of the order of magnitude of atrazine or slightly higher. Among the compounds investigated, only AM is a specific metabolite of atrazine that may indicate human

exposure, but it was never detected in wastewater. The other metabolites detected can also result from exposure to other triazines, particularly terbutylazine, which is the only chlorotriazine herbicide approved for use in EU, and DES, DIA and DEA can originate from degradation of the parent substances in the environment (Barr et al., 2007). It was therefore very difficult to correlate their occurrence in wastewater with human exposure.

The mass loads of pyrethroids were higher than those of triazines, 3-PBA ranged between 4.2 and 30 mg/day/1000 inh and *trans*-DCCA from 7.0 to 46 mg/day/1000 inh. In all the cities, *cis*-DCCA mass loads were the lowest (3.6 - 10.5 mg/day/1000 inh). These specific metabolites were used to evaluate human exposure as described here below.

The sum of the mass loads of the compounds measured for each class of pesticides was calculated as described in paragraph 2.7, in order to compare results from the different cities (Figure 2). Different patterns were observed among the cities and for the various classes of pesticides, but Utrecht and Oslo invariably had the lowest loads. The specific biomarkers of exposure to pyrethroids had the highest loads in Castellon (mean 86 mg/day/1000 inh) followed by Milan and Bristol (mean 43 mg/day/1000 inh), and Copenhagen (mean 41 mg/day/1000 inh). This may indicate a higher human exposure to pyrethroids in Spain due to either direct exposure or consumption of contaminated food, and fits with the fact that Spain is classified as one of the countries with the highest sales of pesticides in Europe (Eurostat, 2014). Regarding the specific metabolites of organophosphates, the highest loads were again in Castellon (mean 28 mg/day/1000 inh), Bristol (mean 26 mg/day/1000 inh) and also in Zurich (mean 21 mg/day/1000 inh). Among non-specific metabolites a direct correlation with exposure could not be performed. The highest levels were found for alkyl phosphates in Zurich (mean 1056 mg/day/1000 inh), followed by Bristol (mean 573 mg/day/1000 inh) and Brussels (mean 322 mg/day/1000 inh), and for triazines in Milan (mean 14 mg/day/1000 inh) Zurich and Brussels (mean 10 mg/day/1000 inh) (Figure 2).

Since human exposure occurs mainly through the diet and can be related to direct exposure only in some cases (i.e. rural areas), the results obtained for the specific biomarkers of exposure can reveal new information about the “average exposure” of the population to these pesticides (pyrethroids and organophosphates). Regarding the other non-specific biomarkers, further investigation will be necessary to assess the main sources of these substances, and exclude the possibility of discharges from sources other than human metabolism.

3.4 Comparison of mass loads of insecticides with official sales statistics

Organophosphates and pyrethroids were the classes most frequently detected in wastewater, both of which are classified as insecticides. Wastewater results were therefore compared with the national sales statistics of insecticides reported by Eurostat (Eurostat, 2014). The sum of the specific biomarkers of insecticides was normalized to the population investigated in each city and the means are reported in Figure 3. Mass loads were the highest in Castellon, Bristol, Copenhagen and Milan and the lowest in Oslo (Figure 3). These results mainly reflect the Eurostat official sales statistics (Figure 3), which reported that Spain, Italy and UK had the highest sales data of insecticides, and Norway had the lowest. Because human exposure to pesticides is mainly influenced by the diet, we can speculate that in the countries with a high sale of insecticides, and a consequent higher use in agriculture, there is also a major supply of products (vegetable and fruits) that leads to a higher exposure to these substances. This is supported by the fact that our study was focused on urban areas where direct exposure related to agricultural use can be excluded. In Spain and Italy the Mediterranean diet, which includes lots of fruits and vegetables, may also play an important role in the exposure to pesticides. Wastewater results seem to reflect also the available figures of vegetable and fruit supply and consumption in Europe which are reported to be higher in the South than in the North of Europe (EUFIC).

397

398 **3.5 Back-calculation of pyrethroid intake**

399 The daily intake by the general population was calculated for pyrethroids due to the
400 suitability of wastewater biomarkers. The mass loads of biomarkers (3-PBA and *trans*- and *cis*-
401 DCCA) were therefore used to back-calculate the intake of the corresponding parent substances.
402 The mass loads of 3-PBA, which is the common urinary metabolic product of about 20 pyrethroids,
403 were multiplied by its specific CF as previously described (Rousis et al., 2017). Pyrethroids highest
404 intake was in Castellon (207 mg/day/1000 inh.) followed by Bristol (77 mg/day/1000 inh.) and
405 Milan (75 mg/day/1000 inh.), and the lowest in Oslo (17 mg/day/1000 inh.) (Table 3).

406 The intake of *trans*- and *cis*- permethrin, cypermethrin and cyfluthrin was estimated using
407 the mass loads of their specific metabolites *trans*- and *cis*-DCCA in wastewater and their specific
408 CF (Rousis et al., 2017). Results are reported in Table 3 as the sum of the *cis*- and *trans*- DCCA
409 isomers. The estimated intakes ranged between 227 mg/day/1000 inh in Castellon and 26 in Oslo.
410 Similar intakes were found in UK (126 mg/day/1000 inh), Copenhagen (123 mg/day/1000 inh) and
411 Milan (130 mg/day/1000 inh).

412 The intake profiles from both DCCA and 3-PBA were highest in Castellon and lowest in
413 Oslo, indicating an extremely divergent exposure to this class of pesticides. These results are in
414 accordance with the eEuropean statistics of fruit and vegetable consumption and also with national
415 statistics of pesticides sales as previously discussed for the entire class of insecticides. The intake of
416 pyrethroids estimated from DCCA was generally higher than those estimated from 3-PBA in all the
417 cities (in several cases the difference was statistically significant, DCCA vs. 3-PBA) (Table 3). This
418 may reflect different patterns of exposure to pyrethroids, which are excreted as the investigated
419 biomarkers. Further research is therefore required to investigate the specific patterns of the
420 household use of these substances and the food contamination.

421

422 3.6 Comparison of estimated intake with the acceptable daily intake (ADI)

423 The potential risk related to the intake of permethrin, cypermethrin and cyfluthrin was
424 assessed using the daily intake estimated from the loads of *trans*- and *cis*-DCCA measured in
425 wastewater. In order to compare these data with ADI values, the ADI of beta-cyfluthrin was used as
426 a worst case scenario, since it was the lowest for this class of compounds. An ADI of 0.003 mg/kg
427 body weight per day for a man of 70 kg resulted in an average consumption of 0.21 mg/person per
428 day (Rousis et al., 2017). The comparison between intakes estimated by WBE and the %ADI are
429 reported in Table 4. The estimated intake of permethrin, cypermethrin and cyfluthrin in the
430 population was generally lower than the ADI, and exceeded this reference value only in one case
431 (Castellon) (Table 4). As previously discussed, this area was found to have the highest exposure
432 level to insecticides (particular pyrethroids) probably due to a combination of wide use of
433 pesticides and high consumption of contaminated food.

434 3.7 Limitations and future research needs

435 Up to date we checked the formation of metabolites from the parent substances through
436 laboratory tests performed in wastewater mimicking different temperature conditions during in-
437 sewer transport and sampling. Nevertheless, it would be ideal to perform transformation
438 experiments in real sewers, but many factors make troublesome to obtain accurate results in such
439 studies. Moreover, the stability of biomarkers in wastewater can be highly affected by “local”
440 conditions in a WWTP and may require specific investigations. Future research in this area should
441 take into account the main processes occurring in sewer compartments, and consequently the
442 potential presence of pesticides/metabolites in the different compartments: a) the bulk liquid
443 (wastewater with suspended particulate matter); b) biofilm growing on the sewer walls; c)
444 sediments; d) the sewer atmosphere (McCall et al., 2016).

The present study is the first one in which an attempt is made to correlate the mass loads of insecticides obtained from WBE with national sales statistics and vegetable and fruit consumption. A number of limitations must be considered to improve future comparisons of this kind of data. On one side, WBE results were obtained by measuring a few specific urinary metabolites that indicate the exposure to a limited number of parent substances within the entire class of insecticides. Furthermore, WBE was performed only in one city per country and for a limited period (seven consecutive days). Thus, results may not reflect longer periods of exposure. Under these conditions, the extrapolation of results to the whole country will be biased by the specific spatial and temporal profiles of that city. This was seen in previous studies, where significant differences in pesticide intake were found among cities within the same country (Rousis et al., 2016), and pesticides levels showed seasonal variations (Rousis et al., 2017). Thus, future WBE studies should include more cities per country and sampling should be repeated seasonally to improve the comparability of wastewater results with the available national statistics. On the other side, national sales statistics for pesticides may not reflect the actual use of these substances in a country and they are obviously not directly related to exposure, even if the first results suggest a correlation. Moreover, these data are referred to the sales of an entire class of substances, for instance insecticides in our case, registered in an EU database and collected over the whole year in each country, being therefore more comprehensive and aggregated than our information from WBE. Finally, food consumption can be measured in different ways and statistics can be obtained with different methods which are not directly comparable. Since National Authorities often adopt different methods to collect data, the comparability of international statistics should be carefully verified.

4 Conclusions

WBE was applied here for the first time to assess human exposure to different classes of pesticides across Europe. Several selected biomarkers of exposure to pesticides were measured in

raw wastewater and used as indicators of human exposure in the population. Mass loads suggested a different pattern of exposure to organophosphates, pyrethroids and triazines. Spatial differences in exposure to insecticides in the various cities were in line with national statistics related to pesticides exposure. Results suggested that in the countries with higher insecticides sales, there is also a major supply of products (vegetables and fruits) that leads to a higher exposure to these substances. WBE was able to provide new information about the “average exposure” of the population to pesticides. Moreover, the calculation of the daily intake of pyrethroids highlighted also a different pattern of exposure within this class. The comparison of daily intake calculated for permethrin, cypermethrin and cyfluthrin and a worst case ADI (the one from beta-cyfluthrin) indicated a potential risk for human health. This study suggest that WBE as can be a very promising complementary biomonitoring tool to evaluate population-wide exposure to pesticides. Some current limitations were also discussed in order to improve future applications.

Contributions

Nikolaos I. Rousis, Sara Castiglioni and Ettore Zuccato planned and designed the study. The collection of the wastewater samples was organized by all authors. Nikolaos I. Rousis analyzed the samples and interpreted the results with the input of Emma Gracia-Lor and Sara Castiglioni. Nikolaos I. Rousis and Sara Castiglioni drafted the manuscript, which was critically revised by all co-authors. All authors are aware of the content and accept responsibility, for the manuscript.

Acknowledgements

Nikolaos I. Rousis, Richard Bade, Jose Antonio Baz-Lomba, Erika Castrignanò, Ana Causanilles, Juliet Kinyua, Ann-Kathrin McCall, Pedram Ramin, Yeonsuk Ryu acknowledge the contribution of the European Union's Seventh Framework Programme under Grant Agreement No.

494 [Marie Curie-FP7-PEOPLE Grant #317205 - SEWPROF] for their Early Stage Researcher (ESR)
495 contracts and Emma Gracia-Lor and Zhugen Yang for their Experienced Researcher (ER) contract.
496 Emma Gracia-Lor is also grateful for financial support from Generalitat Valenciana, Conselleria
497 d'Educació, Investigació, Cultura i Esport (APOSTD/2015, Programa VALi+d) for her post-
498 doctoral contract.

499

References

- Allen, M.T., Levy, L.S., 2013. Parkinson's disease and pesticide exposure – a new assessment. *Crit. Rev. Toxicol.* 43, 515–534. doi:10.3109/10408444.2013.798719
- Aprea, M.C., 2012. Environmental and biological monitoring in the estimation of absorbed doses of pesticides. *Toxicol. Lett.* 210, 110–118. doi:10.1016/j.toxlet.2011.08.008
- Banta-Green, C.J., Field, J.A., Chiaia, A.C., Sudakin, D.L., Power, L., de Montigny, L., 2009. The spatial epidemiology of cocaine, methamphetamine and 3,4-methylenedioxymethamphetamine (MDMA) use: a demonstration using a population measure of community drug load derived from municipal wastewater. *Addiction* 104, 1874–1880. doi:10.1111/j.1360-0443.2009.02678.x
- Barr, D.B., 2008. Biomonitoring of exposure to pesticides. *J. Chem. Heal. Saf.* 15, 20–29. doi:10.1016/j.jchas.2008.07.001
- Barr, D.B., Bravo, R., Weerasekera, G., Calabiano, L.M., Whitehead, R.D., Olsson, A., Caudill, S.P., Schober, S.E., Pirkle, J.L., Sampson, E.J., Jackson, R.J., Needham, L.L., 2004. Concentrations of dialkyl phosphate metabolites of organophosphorus pesticides in the U.S. population. *Environ. Health Perspect.* 112, 186–200. doi:10.1289/ehp.6503
- Barr, D.B., Olsson, A.O., Wong, L.-Y., Udunka, S., Baker, S.E., Whitehead, R.D., Magsumbol, M.S., Williams, B.L., Needham, L.L., 2010. Urinary Concentrations of Metabolites of Pyrethroid Insecticides in the General U.S. Population: National Health and Nutrition Examination Survey 1999–2002. *Environ. Health Perspect.* 118, 742–748. doi:10.1289/ehp.0901275
- Barr, D.B., Panuwet, P., Nguyen, J. V., Udunka, S., Needham, L.L., 2007. Assessing Exposure to Atrazine and Its Metabolites using Biomonitoring. *Environ. Health Perspect.* 115, 1474–1478. doi:10.1289/ehp.10141
- Castiglioni, S., Bijlsma, L., Covaci, A., Emke, E., Hernández, F., Reid, M., Ort, C., Thomas, K. V., van Nuijs, A.L.N., de Voogt, P., Zuccato, E., 2013. Evaluation of Uncertainties Associated with the Determination of Community Drug Use through the Measurement of Sewage Drug Biomarkers. *Environ. Sci. Technol.* 47, 1452–1460. doi:10.1021/es302722f
- Castiglioni, S., Thomas, K. V., Kasprzyk-Hordern, B., Vandam, L., Griffiths, P., 2014. Testing wastewater to detect illicit drugs: State of the art, potential and research needs. *Sci. Total*

- 530 Environ. 487, 613–620. doi:10.1016/j.scitotenv.2013.10.034
- 531 Commision, E., 2008. Directive 2008/105/EC of the European Parliament and of the Council of 16
532 December 2008 on environmental quality standards in the field of water policy, amending and
533 subsequently repealing Council Directives 82/176/EEC, 83/513/EEC, 84/156/EEC,
534 84/491/EEC,.
- 535 Commision, E., 2006. Directive 2006/118/EC of the European Parliament and of the Council of 12
536 December 2006 on the protection of groundwater against pollution and deterioration.
- 537 Commision, E., 1998. Council Directive 98/83/EC of 3 November 1998 on the quality of water
538 intended for human consumption.
- 539 Cooper, J., Dobson, H., 2007. The benefits of pesticides to mankind and the environment. Crop
540 Prot. 26, 1337–1348. doi:10.1016/j.cropro.2007.03.022
- 541 Eurostat, 2014. Pesticide sales statistics. URL [http://ec.europa.eu/eurostat/statistics-](http://ec.europa.eu/eurostat/statistics-explained/index.php/Pesticide_sales_statistics)
542 [explained/index.php/Pesticide_sales_statistics](http://ec.europa.eu/eurostat/statistics-explained/index.php/Pesticide_sales_statistics)
- 543 Garrison, A.W., 2004. On the issue of enantioselectivity of chiral pesticides: a green chemistry
544 opportunity. Green Chem. 6, G77. doi:10.1039/b414799p
- 545 Godfray, H.C.J., Beddington, J.R., Crute, I.R., Haddad, L., Lawrence, D., Muir, J.F., Pretty, J.,
546 Robinson, S., Thomas, S.M., Toulmin, C., 2010. Food Security: The Challenge of Feeding 9
547 Billion People. Science 327, 812–818. doi:10.1126/science.1185383
- 548 Gracia-Lor, E., Zuccato, E., Castiglioni, S., 2016. Refining correction factors for back-calculation
549 of illicit drug use. Sci. Total Environ. 573, 1648–1659. doi:10.1016/j.scitotenv.2016.09.179
- 550 Grube, A., Donaldson, D., Timothy Kiely, A., Wu, L., 2011. Pesticides Industry Sales and Usage:
551 2006 and 2007 Market Estimates. U.S. Environ. Prot. Agency.
- 552 Hernández, F., Sancho, J. V, Pozo, O.J., 2002. Direct determination of alkyl phosphates in human
553 urine by liquid chromatography/electrospray tandem mass spectrometry. Rapid Commun.
554 Mass Spectrom. 16, 1766–1773. doi:10.1002/rcm.790
- 555 Heudorf, U., Angerer, J., 2001. Metabolites of Pyrethroid Insecticides in Urine Specimens: Current
556 Exposure in an Urban Population in Germany. Environ. Health Perspect. 109, 213–217.
557 doi:10.1289/ehp.01109213

- 558 López, Ó., Fernández-Bolaños, J.G., Gil, M.V., 2005. New trends in pest control: the search for
559 greener insecticides. *Green Chem.* 7, 431–442. doi:10.1039/b500733j
- 560 Mamane, A., Baldi, I., Tessier, J.-F., Raheison, C., Bouvier, G., 2015. Occupational exposure to
561 pesticides and respiratory health. *Eur. Respir. Rev.* 24, 306–319.
562 doi:10.1183/16000617.00006014
- 563 McCall, A.-K., Bade, R., Kinyua, J., Lai, F.Y., Thai, P.K., Covaci, A., Bijlsma, L., van Nuijs,
564 A.L.N., Ort, C., 2016. Critical review on the stability of illicit drugs in sewers and wastewater
565 samples. *Water Res.* 88, 933–947. doi:10.1016/j.watres.2015.10.040
- 566 McKelvey, W., Jacobson, J.B., Kass, D., Barr, D.B., Davis, M., Calafat, A.M., Aldous, K.M., 2013.
567 Population-Based Biomonitoring of Exposure to Organophosphate and Pyrethroid Pesticides in
568 New York City. *Environ. Health Perspect.* 121, 1349–1356. doi:10.1289/ehp.1206015
- 569 Ort, C., van Nuijs, A.L.N., Berset, J.-D., Bijlsma, L., Castiglioni, S., Covaci, A., de Voogt, P.,
570 Emke, E., Fatta-Kassinos, D., Griffiths, P., Hernández, F., González-Mariño, I., Grabic, R.,
571 Kasprzyk-Hordern, B., Mastroianni, N., Meierjohann, A., Nefau, T., Östman, M., Pico, Y.,
572 Racamonde, I., Reid, M., Slobodnik, J., Terzic, S., Thomaidis, N., Thomas, K. V., 2014.
573 Spatial differences and temporal changes in illicit drug use in Europe quantified by wastewater
574 analysis. *Addiction* 109, 1338–1352. doi:10.1111/add.12570
- 575 Reemtsma, T., Lingott, J., Roegler, S., 2011. Determination of 14 monoalkyl phosphates, dialkyl
576 phosphates and dialkyl thiophosphates by LC-MS/MS in human urinary samples. *Sci. Total*
577 *Environ.* 409, 1990–1993. doi:10.1016/j.scitotenv.2011.01.032
- 578 Rousis, N.I., Zuccato, E., Castiglioni, S., 2016. Monitoring population exposure to pesticides based
579 on liquid chromatography-tandem mass spectrometry measurement of their urinary metabolites
580 in urban wastewater: A novel biomonitoring approach. *Sci. Total Environ.* 571, 1349–1357.
581 doi:10.1016/j.scitotenv.2016.07.036
- 582 Rousis, N.I., Zuccato, E., Castiglioni, S., 2017. Wastewater-based epidemiology to assess human
583 exposure to pyrethroid pesticides. *Environ. Int.* 99, 213–220. doi:10.1016/j.envint.2016.11.020
- 584 Saillenfait, A.-M., Ndiaye, D., Sabaté, J.-P., 2015. Pyrethroids: Exposure and health effects – An
585 update. *Int. J. Hyg. Environ. Health* 218, 281–292. doi:10.1016/j.ijheh.2015.01.002
- 586 Stallones, L., Beseler, C.L., 2016. Assessing the connection between organophosphate pesticide
587 poisoning and mental health: A comparison of neuropsychological symptoms from clinical

observations, animal models and epidemiological studies. *Cortex* 74, 405–416.
doi:10.1016/j.cortex.2015.10.002

Thomas, K. V., Reid, M.J., 2011. What Else Can the Analysis of Sewage for Urinary Biomarkers
Reveal About Communities? *Environ. Sci. Technol.* 45, 7611–7612. doi:10.1021/es202522d

Venkatesan, A. K., Halden R. U., 2014. Wastewater Treatment Plants as Chemical Observatories to
Forecast Ecological and Human Health Risks of Manmade Chemicals. *Sci. Rep.* 4, 3731. doi:
10.1038/srep03731

Wessels, D., Barr, D.B., Mendola, P., 2003. Use of Biomarkers to Indicate Exposure of Children to
Organophosphate Pesticides: Implications for a Longitudinal Study of Children's
Environmental Health. *Environ. Health Perspect.* 111, 1939–1946. doi:10.1289/ehp.6179

Ye, M., Beach, J., Martin, J.W., Senthilselvan, A., 2015. Urinary Dialkyl Phosphate Concentrations
and Lung Function Parameters in Adolescents and Adults: Results from the Canadian Health
Measures Survey. *Environ. Health Perspect.* 124, 491–497. doi:10.1289/ehp.1509745

Yusa, V., Millet, M., Coscolla, C., Pardo, O., Roca, M., 2015. Occurrence of biomarkers of
pesticide exposure in non-invasive human specimens. *Chemosphere* 139, 91–108.
doi:10.1016/j.chemosphere.2015.05.082

Zuccato, E., Chiabrando, C., Castiglioni, S., Bagnati, R., Fanelli, R., 2008. Estimating community
drug abuse by wastewater analysis. *Environ. Health Perspect.* 116, 1027–1032.
doi:10.1289/ehp.11022

608

609 **Table 1.** Summary of the main characteristics of the metabolites selected as WBE biomarkers.

Metabolites selected as WBE biomarkers	Parent pesticides	Detection in wastewater (present study)	Other potential sources (Rousis et al., 2016)	Stability in wastewater (Rousis et al., 2016)	Formation from parent pesticides in wastewater (Rousis et al., 2017); present study)
Triazines					
DES	Terbutylazine	Yes	Yes	Yes	No
DIA	Atrazine, terbutylazine, simazine, propazine	Yes	Yes	Yes	No
DEA	Atrazine, terbutylazine, simazine, propazine	Yes	Yes	Yes	No
AM	Atrazine	No	Yes	Yes	No
Pyrethroids					
3-PBA	20 pyrethroids ^a	Yes	Yes	Yes	No
trans-DCCA	Permethrin, cypermethrin, cyfluthrin	Yes	No	Yes	No
cis-DCCA	Permethrin, cypermethrin, cyfluthrin	Yes	No	Yes	No
Organophosphates					
TCPY	Chlorpyrifos, chlorpyrifos-methyl	Yes	Yes	Yes	No
MMA	Malathion	Yes	Yes	Yes	No
IMPY	Diazinon	Yes	Yes	Yes	No
DEP	Several organophosphate insecticides	Yes	Yes	Yes	- ^b
DETP	Several organophosphate insecticides	Yes	Yes	Yes	- ^b
DMP	Several organophosphate insecticides	Yes	Yes	No	- ^b
DMTP	Several organophosphate insecticides	No	Yes	Yes	- ^b

610 ^aPermethrin, cypermethrin, deltamethrin, fenvalerate, phenothrin, cyphenothrin, cyhalothrin, esfenvalerate,
611 fenpropathrin, allethrin, resmethrin, tralomethrin, flucythrinate, fluvalinate and their isomers; ^b not assessed
612 because these compounds come from multiple substances.
613

614 **Table 2** Mean concentrations (ng/L) and standard deviations of the raw wastewater samples collected in eight European cities in March 2015.

Compound	Bristol	Brussels	Castellon	Copenhagen	Milan	Oslo	Utrecht	Zurich	Frequency of detection (%)
Triazines									
ATZ	4.4 ± 0.4	12.8 ± 1.3	2.0 ± 1.0	1.3 ± 0.1	7.9 ± 0.8	1.7 ± 0.2	2.1 ± 0.3	5.4 ± 0.6	98.2
DES	<0.6.	<0.6.	21.1 ± 3.7	<0.6.	12.2 ± 1.4	<0.6.	<0.6.	6.2 ± 0.8	37.5
DIA	<1.4	6.7 ± 2.0	<1.4	<1.4	8.9 ± 1.4	<1.4	<1.4	4.3 ± 0.2	36.2
DEA	7.5 ± 3.0	19.6 ± 5.5	4.5 ± 1.2	<1.1	7.7 ± 1.1	<1.1	<1.1	7.4 ± 0.9	62.5
AM	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	<0.5	0
Pyrethroids									
3-PBA	49 ± 25	22.4 ± 1.4	129 ± 32	12.4 ± 2.3	26.1 ± 9.3	5.3 ± 1.5	30.1 ± 7.4	9.6 ± 1.4	98.2
<i>trans</i> -DCCA	118 ± 65	65 ± 13	200 ± 60	44 ± 16	63 ± 34	15.1 ± 8.8	124 ± 54	31 ± 10	96.4
<i>cis</i> -DCCA	22 ± 11	<7.7	45 ± 11	<7.7	14 ± 11	<7.7	22.9 ± 8.3	<7.7	35.7
Organophosphates									
CPF	<2.4	<2.4	<2.4	<2.4	<2.4	<2.4	<2.4	<2.4	0
CPF-MET	<3.5	<3.5	<3.5	<3.5	<3.5	<3.5	<3.5	<3.5	0
TCPY	43 ± 23	23.8 ± 2.7	93 ± 23	17.8 ± 2.3	20.1 ± 2.9	8.3 ± 1.3	28.3 ± 3.9	26.4 ± 3.1	100
MMA isomer 1	<3.9	<3.9	397 ± 966	<3.9	4.7 ± 2.3	<3.9	<3.9	<3.9	8.9
MMA isomer 2	<4.8	<4.8	285 ± 661	<4.8	<4.8	<4.8	<4.8	<4.8	7.1
IMPY	72 ± 48	4.9 ± 1.1	25 ± 11	3.6 ± 0.8	<1.29	6.5 ± 1.2	12.7 ± 2.8	19 ± 16	87.5
Alkyl phosphates (Organophosphates)									
DEP	1076 ± 670	180 ± 24	231 ± 56	110 ± 12	123 ± 20	46 ± 19	206 ± 13	187 ± 22	94.6
DETP	39 ± 19	<17.5	<17.5	<17.5	<17.5	<17.5	<17.5	<17.5	7.1
DMP	1388 ± 2228	1072 ± 1018	278 ± 77	280 ± 92	128 ± 22	233 ± 60	269 ± 43	2269 ± 630	100
DMTP	<395	<395	<395	<395	<395	<395	<395	<395	0

625 <LOQ/2 are reported as used for further calculation. LOQ values are reported in Table S2.

626

627 **Table 3** Pyrethroid intake (mg/day/1000 inhabitants; mean and standard deviation) back-calculated
 628 from 3-PBA and *cis*- and *trans*-DCCA.

WWTP	Group of pyrethroids (3-PBA)	Permethrin, cypermethrin and cyfluthrin (DCCA*)	Statistical analysis (p-values) [§]
Bristol	77 ± 37	126 ± 60	0.091
Brussels	41 ± 6	62 ± 11	0.012
Castellon	207 ± 47	227 ± 59	0.507
Copenhagen	57 ± 13	123 ± 50	0.005
Milan	75 ± 39	130 ± 101	0.209
Oslo	17 ± 5	26 ± 13	0.128
Utrecht	33 ± 8	90 ± 36	0.001
Zurich	29 ± 6	50 ± 22	0.031

629 *Sum of *cis*- and *trans*-DCCA;[§] unpaired t-test or Mann-Whitney test were performed considering a
 630 statistical significance for p<0.05.

631

632

633 **Table 4** Estimated intake of permethrin, cypermethrin and cyfluthrin of the population living in
 634 different European cities and comparison with the acceptable daily intake (ADI) for beta-cyfluthrin
 635 (0.21 mg/day/person).

WWTP	Intake of permethrin, cypermethrin and cyfluthrin (mg/day/person)	% ADI*
Bristol	0.126 ± 0.060	60
Brussels	0.062 ± 0.011	30
Castellon	0.227 ± 0.059	108
Copenhagen	0.123 ± 0.050	58
Milan	0.130 ± 0.101	62
Oslo	0.026 ± 0.013	12
Utrecht	0.090 ± 0.036	43
Zurich	0.050 ± 0.022	24

636 *Permethrin, cypermethrin and cyfluthrin intake percentage compared to the ADI of beta-cyfluthrin and
 637 expressed in %.


638


639


640 **Figure Legends**641 **Fig. 1.** Cities investigated in the present study in Europe.

642 **Fig. 2.** Sum of the mass loads (mg/day/1000 inhabitants) of organophosphates, triazines,
643 pyrethroids and alkyl phosphates in eight European cities.

644 **Fig.3.** Sum of the mass loads of insecticides (mg/day/1000 inhabitants) estimated from wastewater
645 in eight European cities and national sales from Eurostat (2014).







Highlights

- WBE was applied for the first time to assess human exposure to pesticides in Europe
- Different patterns were observed among the cities and the classes of pesticides
- Results were in line with the national statistics related to pesticides exposure
- This study gives new information about the “average exposure” of the population